



Eur päisches Patentamt
European Patent Office
Office européen des brevets

Publication number:

**0 386 951
A2**

12

EUROPEAN PATENT APPLICATION

21 Application number: 90302236.6

51 Int. Cl.⁵: **A61K 37/02, A61K 9/08,
A61K 47/04**

22 Date of filing: 02.03.90

30 Priority: 06.03.89 US 319248

43 Date of publication of application:
12.09.90 Bulletin 90/37

84 Designated Contracting States:
AT BE CH DE DK ES FR GB GR IT LI LU NL SE

71 Applicant: **ELI LILLY AND COMPANY**
Lilly Corporate Center
Indianapolis Indiana 46285(US)

72 Inventor: **Inman, Eugene Lee**
8061 Castle Cove Road
Indianapolis, Indiana 46256(US)
Inventor: **Kirsch, Lee Edwin**
7630 Camelback Drive
Indianapolis, Indiana 46250(US)

74 Representative: **Hudson, Christopher Mark et al**
Erl Wood Manor
Windlesham Surrey GU20 6PH(GB)

54 An improved diluent formulation for daptomycin.

57 Improved formulations for the lipopeptide antibiotic, daptomycin, are provided. Also provided are an improved kit for parenteral administration of daptomycin and a buffered pharmaceutical composition comprising daptomycin and a parenterally-acceptable buffer.

EP 0 386 951 A2

The present invention provides a solution to the problem of the premature "in solution" degradation of the known lipopeptide antibiotic daptomycin. Daptomycin is represented by the following structural Formula

[illegible]

First, 8 vials of daptomycin (100 mg each) were reconstituted to a concentration of 1 mg/ml with normal saline and 5% dextrose (4 each). Next, the solutions were stored at room temperature (approximately 25 °C) and refrigerated temperatures and assayed at the 6 h and 24 h timepoints.

Time			Clarity		Potency	
(hrs.)	Temp.	pH	(NTU)	Visual	(mg/mL)	Total Appearance of Degradation Products
0.9% NaCl:						
Initial	--	4.2	1.0	ACPL*	1.026	5.2
6	A	4.2	0.9	1111*	1.025	5.1
	B	4.2	0.8	1111	1.025	5.6
24	A	4.2	0.9	1111	1.019	5.6
	B	4.2	0.7	1111	1.002	7.5
5% Dextrose:						
Initial	--	4.5	1.7	ACPL	1.023	8.9
6	A	4.5	1.6	1111	1.001	12.6
	B	4.5	0.7	1111	0.963	16.6
24	A	4.5	1.5	1111	0.975	16.2
	B	4.5	1.1	1111	0.848	24.9

*Inspection for appearance, color, package and clarity.

*1 indicates no change from initial.

In sum, solutions of daptomycin and dextrose held at refrigerated temperature demonstrated a potency loss of about 5% at the 24 h timepoint, while solutions held at room temperature showed a 17% loss over the same time period.

Further, a series of degradation kinetic studies demonstrate that the rate of appearance of degradation products is pH-dependent. For example, the appearance half-life for the degradation products at 25 °C and pH 4.5 was 6 h, while at 25 °C and pH 7.0, the half-life was 63 h.

The nature of the degradation products has as yet not been determined; although, as indicated by the loss in potency of unstabilized solutions of daptomycin, these products exhibit far less antibiotic activity than the present antibiotic.

The stabilized formulations provided herein are suitable for administration by the parenteral routes and comprise an antibiologically effective amount of daptomycin, a pharmaceutically acceptable buffer and a pharmaceutically acceptable diluent. In addition, the formulations can contain excipients.

Pharmaceutically acceptable buffers which can be used include sodium or potassium salts of phosphoric acid, boric acid, citric acid, carbonic acid, hydroxide, and the like; tris(hydroxymethyl)aminomethane (TRIS® buffer), amino acids, and like buffers.

Diluents which can be used include water for injection, 5% dextrose, 0.9% saline, Ringer's solution and like commonly used diluents for the parenterally administered antibiotics.

Excipients which can be used include, for example, mannitol, tonicity modifiers (e.g., sodium chloride or glycerol), preservatives and solubilizing agents. Further examples of acceptable diluents and excipients can be found in Remington's Pharmaceutical Sciences, 17th Ed., A. R. Gennaro, Ed., Mack Publishing Co., Easton, PA, 1985, incorporated herein by reference; particularly relevant are pages 1518-1552.

Preferred buffers of this invention are dibasic sodium phosphate and TRIS® (tris(hydroxymethyl)-aminomethane buffer).

Thus, in broadest terms, the present invention provides an improved formulation comprising daptomycin and a parenterally-acceptable buffer possessing sufficient buffer capacity to maintain the pH of the reconstituted daptomycin between a pH of between about 6.0 to about 8.0.

Specific examples of preferred parenterally acceptable buffer systems are provided in the following Table 1:

Table 1

Formulation A	Isotonic Sodium Phosphate Dibasic Solution (8.73 mg/mL)
Formulation B	Isotonic Sodium Phosphate Dibasic Solution (15.76 mg/mL)
Formulation C	TRIS® (Tris(hydroxymethyl)aminomethane) Solution (3.30 mg/mL)
Formulation D	TRIS® (Tris(hydroxymethyl)aminomethane) Solution (3.74 mg/mL)
Formulation E	TRIS® (Tris(hydroxymethyl)aminomethane) Solution (4.96 mg/mL)

Formulations comprising the buffers in Table 1 were tested by reconstituting freeze-dried daptomycin (150 mg) and 50 mg mannitol (50 mg) in 10 mL of each buffer, then diluting the resultant solution in 100 mL of 5% dextrose solutions. The effectiveness of the buffered diluents in suppressing the appearance of degradation products is shown below in Table 2:

Table 2:

Summary Stability Results . Percent Increase in Daptomycin Degradation Products at 1 day storage (25 ° C) and 7 days storage (5 ° C).			
		% Increase in Dextrose Related Degradation Products	
Formulation	pH	1 day (25 ° C)	7 days (5 ° C)
5% dextrose (control)	4.4	16.9	17.3
A	6.9	1.1	1.1
B	7.3	0.6	0.5
C	6.9	1.8	0.8
D	7.4	0.5	0.4
E	7.9	0.3	0.3

Further examples of buffer systems meant by the term "pharmaceutically-acceptable buffer" include sodium or potassium salts of phosphoric acid, boric acid, citric acid, carbonic acid, hydroxide, and the like; tris(hydroxymethyl)aminomethane (TRIS® buffer) and all amino acids and like buffers.

In the case where sodium phosphate dibasic is used as buffer, it is preferred that the concentration be in the range of about 0.02 mMolar to about 0.13 mMolar, most preferably about 0.05 mMolar to about 0.10 mMolar. One skilled in the art will appreciate that the amount of buffer necessary to maintain the pH in the desired range is directly related to the amount of daptomycin present in solution. Thus, the above ranges are preferred when daptomycin is present in a concentration of about 6×10^{-4} mMolar.

The following is a nonexhaustive list of examples of unit dose formulations for daptomycin:

	<u>Ingredient</u>	<u>Amount</u>
5	1) Daptomycin	150 mg
	Mannitol	50 mg
	Isotonic Sodium Phosphate	10 ml
10	Dibasic Solution (8.73 mg/ml)	
	Dextrose (5%)	90 ml
15	2) Daptomycin	200 mg
	Mannitol	66 mg
	Isotonic Sodium Phosphate	13 ml
	Dibasic Solution (8.73 mg/ml)	
20	Dextrose (5%)	87 ml
25	3) Daptomycin	250 mg
	Mannitol	82 mg
	Isotonic Sodium Phosphate	16 ml
	Dibasic Solution (8.73 mg/ml)	
30	Dextrose (5%)	84 ml
35	4) Daptomycin	300 mg
	Mannitol	100 mg
	Isotonic Sodium Phosphate	20 ml
	Dibasic Solution (8.73 mg/ml)	
40	Dextrose (5%)	80 ml

40

45

50

55

5	5)	Daptomycin	150 mg
		Mannitol	50 mg
		Isotonic Sodium Phosphate	10 ml
		Dibasic Solution (15.76 mg/ml)	
		Dextrose (5%)	90 ml
10	6)	Daptomycin	200 mg
		Mannitol	60 mg
		Isotonic Sodium Phosphate	13 ml
15		Dibasic Solution (15.76 mg/ml)	
		Dextrose (5%)	87 ml
20	7)	Daptomycin	250 mg
		Mannitol	82 mg
		Isotonic Sodium Phosphate	16 ml
		Dibasic Solution (15.76 mg/ml)	
25		Dextrose (5%)	84 ml
30	8)	Daptomycin	300 mg
		Mannitol	100 mg
		Isotonic Sodium Phosphate	20 ml
		Dibasic Solution (15.76 mg/ml)	
35		Dextrose (5%)	80 ml
40	9)	Daptomycin	150 mg
		Mannitol	50 mg
		Tris(hydroxymethyl)amminomethane	10 ml
		Solution (3.30 mg/ml)	
45		Dextrose (5%)	90 ml

5	10)	Daptomycin	200 mg
		Mannitol	66 mg
		Tris(hydroxymethyl)amminomethane	13 ml
		Solution (3.30 mg/ml)	
		Dextrose (5%)	87 ml
10	11)	Daptomycin	250 mg
		Mannitol	82 mg
		Tris(hydroxymethyl)amminomethane	16 ml
15		Solution (3.30 mg/ml)	
		Dextrose (5%)	84 ml
20	12)	Daptomycin	300 mg
		Mannitol	100 mg
		Tris(hydroxymethyl)amminomethane	20 ml
		Solution (3.30 mg/ml)	
25		Dextrose (5%)	80 ml
30	13)	Daptomycin	150 mg
		Mannitol	50 mg
		Tris(hydroxymethyl)amminomethane	10 ml
		Solution (3.74 mg/ml)	
35		Dextrose (5%)	90 ml
40	14)	Daptomycin	200 mg
		Mannitol	66 mg
		Tris(hydroxymethyl)amminomethane	13 ml
		Solution (3.74 mg/ml)	
45		Dextrose (5%)	87 ml

5	15)	Daptomycin	250 mg
		Mannitol	82 mg
		Tris(hydroxymethyl)amminomethane	16 ml
		Solution (3.74 mg/ml)	
		Dextrose (5%)	84 ml
10	16)	Daptomycin	300 mg
		Mannitol	100 mg
		Tris(hydroxymethyl)amminomethane	20 ml
15		Solution (3.74 mg/ml)	
		Dextrose (5%)	80 ml
20	17)	Daptomycin	150 mg
		Mannitol	50 mg
		Tris(hydroxymethyl)amminomethane	10 ml
		Solution (4.96 mg/ml)	
25		Dextrose (5%)	90 ml
30	18)	Daptomycin	200 mg
		Mannitol	66 mg
		Tris(hydroxymethyl)amminomethane	13 ml
		Solution (4.96 mg/ml)	
35		Dextrose (5%)	87 ml
40	19)	Daptomycin	250 mg
		Mannitol	82 mg
		Tris(hydroxymethyl)amminomethane	16 ml
		Solution (4.96 mg/ml)	
45		Dextrose (5%)	84 ml
50	20)	Daptomycin	300 mg
		Mannitol	100 mg
		Tris(hydroxymethyl)amminomethane	20 ml
		Solution (4.96 mg/ml)	
55		Dextrose (5%)	80 ml

This aspect of the present invention may be practiced in two ways. First, the daptomycin may be purchased in a lyophilized form and reconstituted prior to use with a buffered diluent as discussed above.

Alternatively, daptomycin may be packaged with a suitable amount of buffer so that upon reconstitution with a (non-buffered) diluent, the resulting reconstituted formulation will also have a pH in the range of about 6 to about 8.

As another aspect of the present invention, there is provided a pharmaceutical composition comprising an antibiotically-effective amount of the antibiotic daptomycin, combined with a parenterally-acceptable buffer in sufficient quantity so as to afford a solution pH of about 6.0 to about 8.0 when diluted with one or more pharmaceutically-acceptable diluents and excipients.

As a further aspect of the present invention, there is provided a parenteral formulation of daptomycin, which comprises admixing daptomycin with one or more pharmaceutically-acceptable diluents and excipients characterized in that a sufficient quantity of a parenterally-acceptable buffer is added to maintain the formulation at a pH of between about 6.0 and about 8.0.

Claims

15

1. A parenteral formulation which comprises as an active ingredient daptomycin associated with a sufficient amount of a parenterally-acceptable buffer to maintain the pH between about 6.0 and about 8.0, and one or more pharmaceutically-acceptable diluents or excipients.

2. A formulation as claimed in claim 1, wherein the buffer is sodium phosphate dibasic.

20

3. A formulation as claimed in claim 1, wherein the buffer is tris(hydroxymethyl)aminomethane.

4. A formulation as claimed in claim 2, wherein the buffer sodium phosphate dibasic is present in the concentration of about 0.02 to about 0.13 mMolar.

5. A formulation as claimed in claim 3, wherein the buffer tris(hydroxymethyl)aminomethane is present in a concentration of about 0.015 to 0.03 mMolar.

25

6. A formulation as claimed in claim 4, which comprises

(a) Daptomycin	150 mg
Mannitol	50 mg
Isotonic Sodium Phosphate	10 ml
Dibasic Solution (8.73 mg/ml)	
Dextrose (5%)	90 ml;

35

40

45

50

55

	(b)	Daptomycin	200 mg
		Mannitol	66 mg
5		Isotonic Sodium Phosphate	13 ml
		Dibasic Solution (8.73 mg/ml)	
		Dextrose (5%)	87 ml;
10	(c)	Daptomycin	250 mg
		Mannitol	82 mg
		Isotonic Sodium Phosphate	16 ml
15		Dibasic Solution (8.73 mg/ml)	
		Dextrose (5%)	84 ml;
20	(d)	Daptomycin	300 mg
		Mannitol	100 mg
		Isotonic Sodium Phosphate	20 ml
		Dibasic Solution (8.73 mg/ml)	
25		Dextrose (5%)	80 ml;
30	(e)	Daptomycin	150 mg
		Mannitol	50 mg
		Isotonic Sodium Phosphate	10 ml
		Dibasic Solution (15.76 mg/ml)	
35		Dextrose (5%)	90 ml;

40

45

50

55

5	(f)	Daptomycin	200 mg
		Mannitol	60 mg
		Isotonic Sodium Phosphate	13 ml
		Dibasic Solution (15.76 mg/ml)	
		Dextrose (5%)	87 ml;
10	(g)	Daptomycin	250 mg
		Mannitol	82 mg
		Isotonic Sodium Phosphate	16 ml
15		Dibasic Solution (15.76 mg/ml)	
		Dextrose (5%)	84 ml;
20	(h)	Daptomycin	300 mg
		Mannitol	100 mg
		Isotonic Sodium Phosphate	20 ml
		Dibasic Solution (15.76 mg/ml)	
25		Dextrose (5%)	80 ml;
30	(i)	Daptomycin	150 mg
		Mannitol	50 mg
		Tris(hydroxymethyl)aminomethane	10 ml
		Solution (3.30 mg/ml)	
35		Dextrose (5%)	90 ml;

40

45

50

55

5	(j)	Daptomycin	200 mg
		Mannitol	66 mg
		Tris(hydroxymethyl)aminomethane	13 ml
		Solution (3.30 mg/ml)	
		Dextrose (5%)	87 ml;
10	(k)	Daptomycin	250 mg
		Mannitol	82 mg
		Tris(hydroxymethyl)aminomethane	16 ml
15		Solution (3.30 mg/ml)	
		Dextrose (5%)	84 ml;
20	(l)	Daptomycin	300 mg
		Mannitol	100 mg
		Tris(hydroxymethyl)aminomethane	20 ml
		Solution (3.30 mg/ml)	
25		Dextrose (5%)	80 ml;
30	(m)	Daptomycin	150 mg
		Mannitol	50 mg
		Tris(hydroxymethyl)aminomethane	10 ml
		Solution (3.74 mg/ml)	
35		Dextrose (5%)	90 ml;

40

45

50

55

	(n)	Daptomycin	200 mg
		Mannitol	66 mg
5		Tris(hydroxymethyl)aminomethane Solution (3.74 mg/ml)	13 ml
		Dextrose (5%)	87 ml;
10	(o)	Daptomycin	250 mg
		Mannitol	82 mg
		Tris(hydroxymethyl)aminomethane Solution (3.74 mg/ml)	16 ml
15		Dextrose (5%)	84 ml;
20	(p)	Daptomycin	300 mg
		Mannitol	100 mg
		Tris(hydroxymethyl)aminomethane Solution (3.74 mg/ml)	20 ml
25		Dextrose (5%)	80 ml;
30	(q)	Daptomycin	150 mg
		Mannitol	50 mg
		Tris(hydroxymethyl)aminomethane Solution (4.96 mg/ml)	10 ml
35		Dextrose (5%)	90 ml;

40

45

50

55

	(r)	Daptomycin	200 mg
		Mannitol	66 mg
5		Tris(hydroxymethyl)aminomethane	13 ml
		Solution (4.96 mg/ml)	
		Dextrose (5%)	87 ml;
10		(s) Daptomycin	250 mg
		Mannitol	82 mg
		Tris(hydroxymethyl)aminomethane	16 ml
15		Solution (4.96 mg/ml)	
		Dextrose (5%)	84 ml; or
		(t) Daptomycin	300 mg
20		Mannitol	100 mg
		Tris(hydroxymethyl)aminomethane	20 ml
		Solution (4.96 mg/ml)	
25		Dextrose (5%)	80 ml.

7. A therapeutic kit for the preparation of a parenteral formulation of the antibiotic daptomycin, said kit comprising at least two containers, one of which contains daptomycin; and a second of which contains a buffer capable of maintaining the pH of a solution of daptomycin in the first container between a pH of about 6.0 and about 8.0 when mixed, each of said first and second containers comprising one or more pharmaceutically acceptable carriers or excipients as appropriate.

8. A pharmaceutical composition comprising an antibiotically-effective amount of the antibiotic daptomycin, combined with a parenterally-acceptable buffer in sufficient quantity so as to afford a solution pH of about 6.0 to about 8.0 when diluted with one or more pharmaceutically-acceptable diluents and excipients.

9. A process for preparing a parenteral formulation of daptomycin, which comprises admixing daptomycin with one or more pharmaceutically-acceptable diluents or excipients, characterized in that a sufficient quantity of a parenterally-acceptable buffer is added to maintain the formulation at a pH of between about 6.0 and about 8.0.